

# Quantum Number Fractionization in Cuprates: U(1) RVB Gauge Theory and Senthil Fisher Constructions

G. Baskaran  
Institute of Mathematical Sciences  
C I T Campus  
Madras 600 113, India

We critically look at Senthil Fisher constructions of quantum number fractionization in cuprates. The first construction, while mathematically correct is of limited relevance to cuprates. The second one has a missing U(1) symmetry. Once this aspect is repaired, it reveals itself as the U(1) RVB gauge theory construction of quantum number fractionization. An approximate local  $Z_2$  symmetry arises in the spin sector for a reason very different from Senthil and Fisher's. The charge sector continues to have the local U(1) symmetry and the metallic spin gap phase of the 2d cuprates may, in principle, carry (irrational) fractional charge  $e^* \neq e$  obeying Haldane's fractional exclusion statistics like the 1d models.

In the RVB theory of cuprates [1,2] the (s)slave-particle formalism [3] and the U(1) gauge theory development [4–6] suggested a natural mechanism of quantum number fractionization in undoped and doped cuprates. The spin charge separation [7] simply corresponded to the deconfinement [4] of s-particles carrying a new U(1) charge called RVB charge [6]. As the theory involves matter (a fermi sea of spinons and bosonic holons) interacting with dynamically generated U(1) gauge fields, spinon deconfinement turned out to be a natural (the only ?) possibility [6] even in the 1 and 2 space dimensional t-J models (in the absence of matter the U(1) gauge theory in 1 and 2 space dimensions have only a confined phase). Soon after the formulation of the RVB gauge theory, a reduction of the local U(1) symmetry to a local  $Z_2$  symmetry was found [8] leading to a theory of  $Z_2$  gauge fields coupled to spinon and holon matter utilizing an important identity due to Marston [9]. This approach has remained unexplored. Several authors, Patrick Lee [10] and collaborators in particular have pursued the gauge theory approach in the last one decade.

Recently Senthil and Fisher (SF) have advocated a construction of quantum number fractionization (QNF)  $2e \rightarrow e + e$  in two different ways [11,12]. In the most recent approach [12], SF study the t-J model in the s-particle formalism and split a neutral d-wave spinon pair into two and use it to construct a spinon and a 'chargin'. We critically analyze this construction [12] and find a missing local U(1) gauge invariance. This invariance is necessary to ensure the crucial double occupancy constraint. We have found a way of correcting this serious difficulty. Once this is done, SF's construction becomes the same as the U(1) RVB gauge theory construction of QNF. In the earlier approach [11], a charge 'e' excitation of the charge '2e' fluid in the metallic spin gap phase of the cuprate is invoked. While this in principle is an allowed possibility in the cuprates we will argue that this is only a particular case allowed by the U(1) RVB gauge theory of the t-J model; a general possibility allowed by the U(1) gauge theory is more consistent with the 'charge

gaplessness' of the metallic spin (pseudo)gap phase.

In the process we also bring out a  $Z_2$  aspect of the RVB gauge theory that was discussed earlier in the game [8,9]. A fundamental difference of our  $Z_2$  gauge symmetry, which is restricted to the spin sector of the doped Mott insulator, from that of SF is also brought out. Having established the connection of SF's construction to the old U(1) RVB mechanism, we also point out how the local U(1) symmetry of the charge sector (in the metallic spin gap phase) provides a more general QNF and the possibility of fractional charge for holons  $e^* \neq e$ . In this way we also suggest that the physics of the charge sector is fairly dimension independent. We also point out that in a (gapless) metallic 2d state such as the spin gap phase with extended (in general having power law form factor) spinons and holons it is more meaningful to talk in terms of the Haldane's exclusion statistics [13] rather than limit oneself to ideas derived from exchange statistics.

First we elaborate how the U(1) RVB gauge theory allows for a unique 'U(1) string' representation of the physical spinon and holon, allowing for a general possibility of charge  $e^* \neq e$  for the holon. Then show how the recent construction of SF, when corrected, becomes the RVB theory's U(1) string representation.

In the (s)slave-particle representation of the the t-J model an electron is decomposed as

$$c_{i\sigma}^\dagger \equiv e_i s_{i\sigma}^\dagger \quad (1)$$

in terms of the 'bare spinon' and 'bare holon' operators  $(s_{i\sigma}^\dagger, e_i^\dagger)$ . The double occupancy constraint takes the form  $\sum_\sigma s_{i\sigma}^\dagger s_{i\sigma} + e_i^\dagger e_i = 1$ . All physical objects are invariant under the local U(1) gauge transformation applied to the s-particles:

$$(s_{i\sigma}^\dagger, e_i^\dagger) \rightarrow e^{-i\theta_i} (s_{i\sigma}^\dagger, e_i^\dagger) \quad (2)$$

In view of the above transformation property, we say that a bare spinon or a holon carry an unit of U(1) 'charge' which we called RVB charge [6]. These constituent s-particles are not physical particles. Physical particles

must remain invariant under the local U(1) gauge transformation. Gauge theory suggests that we attach electric field lines to ‘electric charges’ to make them gauge invariant. In our system the field lines that go to infinity (boundary of the sample) can be created using spin singlet ‘U(1)string’ operators such as:

$$U_{i_1} U_{i_1 i_2} U_{i_2 i_3} \dots \equiv U_l(C) \quad (3)$$

The link variables  $U_{ij}$  are chosen from the three types of spin singlet operators:  $U_{ij} = b_{ij}^\dagger = \frac{1}{\sqrt{2}}(s_{i\uparrow}^\dagger s_{j\downarrow}^\dagger - s_{i\downarrow}^\dagger s_{j\uparrow}^\dagger)$ , or  $\tau_{ij} = \sum_\sigma s_{i\sigma}^\dagger s_{j\sigma}$  or  $e_i^\dagger e_j$  and their complex conjugates. A simple example of the string operator used in ref [4,6] for the Mott insulator is  $b_{l i_1} b_{i_1 i_2}^\dagger b_{i_2 i_3} \dots$ . By construction, under the local U(1) transformation a U(1) string operator transforms as  $U_{l i_1} U_{i_1 i_2} U_{i_2 i_3} \dots \rightarrow e^{i\theta_l} U_{l i_1} U_{i_1 i_2} U_{i_2 i_3} \dots$ . The field line operators represent the ‘spin singlet’ disturbance and charge disturbance created in the spin liquid state in the process of QNF, of transporting a partner spin- $\frac{1}{2}$  moment to the boundary of the system along the sequence C (not necessarily formed by nearest neighbor lattice sites). In a continuum approximation of the U(1) RVB gauge theory the string operators correspond to:  $e^{i \int_{\mathbf{r}} \mathbf{A}_{rvb} \cdot d\mathbf{l}}$ . Here  $\mathbf{A}_{rvb}$  is the dynamically generated U(1) RVB gauge field. That is, schematically a spin one operator or an electron operator undergo the following change during quantum number fractionization:

$$\begin{aligned} S_l^+ &= s_{l\uparrow}^\dagger s_{l\downarrow} \rightarrow s_{l\uparrow}^\dagger U_{l i_1} U_{i_1 i_2} U_{i_2 i_3} \dots U_{i_n \infty} s_{\infty \downarrow} \\ c_{l\sigma}^\dagger &= e_l s_{l\sigma}^\dagger \rightarrow e_l U_{l i_1} U_{i_1 i_2} U_{i_2 i_3} \dots U_{i_n \infty} s_{\infty \sigma}^\dagger \end{aligned}$$

The field line operators in combination with the bare s-particles have the full local U(1) gauge invariance and provides the *most general* localized physical spinon or holon operator characterized by a string C and site l:

$$\zeta_{l\sigma}^{s\dagger} = s_{l\sigma}^\dagger U_l(C) \quad \text{and} \quad \zeta_l^{h\dagger} = e_l^\dagger U_l(C) \quad (4)$$

A propagating low energy spinon or holon eigen state of a spin liquid state exhibiting quantum number fractionization (deconfinement) can be created by the operators:

$$\begin{aligned} \zeta_\sigma^{s\dagger}(\mathbf{q}) &\equiv \sum_{l,C} e^{i\mathbf{q} \cdot \mathbf{r}_l} \phi^s(C, l; \mathbf{q}) s_{l\sigma}^\dagger U_l(C) \\ &\equiv \sum_l e^{i\mathbf{q} \cdot \mathbf{r}_l} s_{l\sigma}^\dagger \eta_l^s(\mathbf{q}) \quad \text{and} \\ \zeta^{h\dagger}(\mathbf{q}) &\equiv \sum_{l,C} e^{i\mathbf{q} \cdot \mathbf{r}_l} \phi^h(C, l; \mathbf{q}) e_l^\dagger U_l(C) \\ &\equiv \sum_l e^{i\mathbf{q} \cdot \mathbf{r}_l} e_l^\dagger \eta_l^h(\mathbf{q}) \end{aligned} \quad (5)$$

The string amplitudes  $\phi^s(C, l; \mathbf{q})$  and  $\phi^h(C, l; \mathbf{q})$  depends on the form of the Hamiltonian and in general may contain some further quantum numbers such as ‘chirality’ for the spinon and holon.

The operator  $\eta_l^{s,h} = \sum_C \phi^{s,h}(C, l; \mathbf{q}) U_l(C)$  creates a quantum delocalized U(1) string excitation starting from a site l; in a classical picture it is a radially flowing U(1) field lines emanating from a site l. We call  $\eta_l^s$  and  $\eta_l^h$  as the ‘radial field operators’ (RFO) of the deconfined phase for the spinon and holon. By construction the RFO’s transform as  $\eta_l \rightarrow e^{i\theta_l} \eta_l$  under the local U(1) gauge transformation.

In a deconfined state the RFO represents the ‘radial’ fashion in which QNF takes place leading to the possibility of a finite energy spinon or holon. The local U(1) gauge invariant physical spinon operator at a site l,  $s_{l\sigma}^\dagger \eta_l^s$  and the holon operator  $e_l^\dagger \eta_l^h$  are the analogue of the finite energy Laughlin quasi hole operator. They create a spin half moment or a missing / excess charge in an otherwise singlet environment, by pushing another spin half moment to the boundary.

In a gapless deconfined state the spin and charge disturbance created (by the RFO), in the process of QNF, will extend to infinity in a power law fashion. We will call this as a power law form factor of the ‘extended’ spinon or holon. The form factors will in general be different for a spinon and a holon. In the 1d Heisenberg and Hubbard models there is a domain wall of spin disturbance of infinite size, around the unpaired spin, as the energy of the spinon tends to zero.

Also the spin SU(2) symmetry (present for zero magnetic field) allows only half integer value of the spin after the quantum number fractionization. In terms of RFO the virtual excitations they create are always of singlet type. Once we have an external magnetic field the spin rotation symmetry is reduced to U(1) symmetry and the spin of the spinon can be different from half, as is also known to be the case for the 1d Heisenberg model. In the presence of the magnetic field the RFO will contain some triplet components, which may change the effective spin of the spinon from half.

The case of holon is more complex in the presence of gapless charge excitation, as in the spin gap metallic phase. The U(1) symmetry associated with the electric charge in general allows any fraction  $e^* \neq e$  for the holon excitations. This is formally achieved by the appropriate choice of the ‘pure’ charge fluctuation terms  $e_{i_s}^\dagger e_{i_{s+1}}$  in the U(1) string, which can push or attract a fractional average charge away from the site l. This is indeed what happens in the 1d Hubbard model. In general, in view of the charge gaplessness of the metallic phase of the 2d cuprates, the charge of the holon should continuously changes as a function of doping x from  $e^* = e$  for the Mott insulator. I do not find a basic principle or a topological reason in the metallic phase of the cuprates which restricts the charge  $e^*$  to e. The power law accumulation of charges and the associated pushing of states to the boundary also leads to the natural possibility of Haldane’s fractional exclusion statistics.

All our discussion was for an infinite system. If we have a finite and large system, a spinon or a holon can not be created in isolation, they have to be created in pairs. Our formal construction can be carried out without any difficulty in this case as well. The above discussion does not prove the existence of deconfined phase in some 2d models, but gives a unique construction for objects with fractional quantum numbers in the s-particle formalism. It should be remarked that the RVB construction is dimension independent, suggesting the likely possibility of a fairly universal character of the quantum number fractionization in the t-J model.

Having discussed the essential aspects in the U(1) gauge mechanism of quantum number fractionization in some detail we go to the construction of SF. Central to their construction [12], for the t-J model in the s-particle representation is a neutral d-wave spinon pair operator  $d_i$  and ‘half’ of the pair  $b_i$  defined in terms of the nearest neighbor valence bond operator  $b_{ij}^\dagger = \frac{1}{\sqrt{2}}(s_{i\uparrow}^\dagger s_{j\downarrow}^\dagger - s_{i\downarrow}^\dagger s_{j\uparrow}^\dagger)$ ,

$$d_i^\dagger \sim \sum_j \alpha_{ij} b_{ij}^\dagger \quad \text{and} \quad (b_i^\dagger)^2 \sim d_i^\dagger \quad (6)$$

And  $\alpha_{ij} = 1$  and  $-1$  for the horizontal and vertical nearest neighbor bonds respectively and zero otherwise. The half of spinon pair  $b_i$  is used to define the ‘spinon’ and ‘chargon’ operators ( $f_{i\sigma}, h_i$ ) through the following relations:

$$e_i \equiv b_i h_i \quad \text{and} \quad s_{i\sigma} \equiv b_i f_{i\sigma} \quad (7)$$

If  $b-i$  transforms as  $b_i \rightarrow e^{i\theta_i} b_i$ , then  $(h_i, f_{i\sigma})$  are gauge invariant physical observables. Let us check how  $b_i$  transforms. On applying the U(1) gauge transformation (equation 2) the  $d_i$  operator transforms as:

$$d_i \rightarrow \sum_j \alpha_{ij} e^{i\theta_i} b_{ij} e^{i\theta_j} = e^{i\theta_i} \sum_j \alpha_{ij} b_{ij} e^{i\theta_j} \quad (8)$$

The local phase factors of the U(1) transformation do not completely factor out. *That is, the d-symmetric spinon pair, as opposed to a valence bond pair, does not carry a definite U(1) charge of the local symmetry.* Consequently, the half of spinon pair  $b_i$  also does not carry unit U(1) charge. Thus the spinon and chargon defined through equation (4) are not neutral under the local U(1) gauge transformation, making them unqualified as physical excitations. One may invoke a coarse graining argument and suggest that if the local U(1) gauge transformation is smooth over the scale of the lattice parameter,  $d_i$  transforms as  $d_i \rightarrow e^{2i\theta_i}$ . However, a coarse graining of a local symmetry is not allowed, as the double occupancy constraint on individual sites is ensured only by the local U(1) invariance.

One may be tempted to split the bond singlet operator as  $b_{ii+\Delta} \sim b_i b_{i+\Delta}$ , to extract an RFO. This is dangerous

and such a split representation for the bond operator fails to reproduce the important commutation relation:

$$[b_{ij}^\dagger b_{ij}, b_{jk}^\dagger b_{jk}] = i \mathbf{S}_j \cdot (\mathbf{S}_i \times \mathbf{S}_k) \quad (9)$$

Is there a way of rectifying the U(1) gauge invariance problem? The answer is obvious once we recognize that the ‘half d-wave spinon pair’ operator  $b_i$  is invented to do the same job that the RFO  $\eta_i^s$  and  $\eta_i^h$  did in the RVB construction. The only objects that carry a U(1) charge (apart from the s-particles themselves) are the open U(1) strings or suitable linear combinations. Open U(1) strings alone can naturalize the U(1) RVB charge of  $s_{i\sigma}^\dagger$  or  $e_i^\dagger$ . The  $Z_2$  string of SF, for the t-J model is a consequence of their erroneous construction, and it does not solve our U(1) problem.

It is also clear that once we extract  $\eta_i$  in a gauge invariant fashion as detailed above, there is no new  $Z_2$  symmetry - we have the full U(1) symmetry. However, for a reason very different from Senthil and Fisher’s, we will show below an approximate reduction of the original U(1) to a  $Z_2$  symmetry arising from the s-particle construction. This is related to the ‘RVB flux’ or ‘RVB magnetic field’ that came up in a concrete fashion from the discovery of Affleck-Marston’s [14]  $\pi$ -flux phase of the Heisenberg antiferromagnet. Later works [15] have also introduced a related  $Z_2$  symmetry in frustrated spin systems. This  $Z_2$  symmetry is also related to the topological order that Kivelson [16] and collaborators were trying to bring out through their study of quantum dimer models and the phase string in Sutherland’s analysis [17] of the short range RVB state.

To understand this in detail in our context, let us go back to the early work by Yu Lu, Tosatti [8] and the present author where the local U(1) gauge theory was reduced to a local  $Z_2$  gauge theory using an important observation of Marston [9]. Consider a closed string made of  $\tau_{ij} = \sum_\sigma s_{i\sigma}^\dagger s_{j\sigma}$ , forming an elementary plaquette:

$$W_{ijkl} \equiv \tau_{ij} \tau_{jk} \tau_{kl} \tau_{li} \sim e^{i \oint \mathbf{A}_{rvb} \cdot d\mathbf{l}} \quad (10)$$

Marston considered the Heisenberg model and found that this plaquette operator satisfies the following important identity in the physical Hilbert space:

$$W_{ijkl}^3 = W_{ijkl} \quad (11)$$

As the plaquette loop  $W_{ijkl}$  is the analogue of the Bohm-Aharonov phase factor of the RVB magnetic field, Marston concluded that the important RVB fluxes enclosed by the plaquette operator are  $Z_2$  in character (0 or  $\pi$  flux, modulo  $2\pi$ ) and he implemented it through a Chern-Simon term in the U(1) gauge theory. On the other hand the authors of reference [8] used Marston identity in an approximate and physical way and reduced the gauge theory from U(1) to  $Z_2$ . *Since Marston identity involves only the spinon s-particles, the reduction of*

$U(1)$  to  $Z_2$  is restricted to the spin sector. The charge sector continues to have the  $U(1)$  symmetry, but is influenced by the condensation of the  $\pi$  flux in the spin sector as shown in ref [8].

We discuss an important limitation in Fisher et al.'s approach in the context of cuprates. They are strongly influenced by the notion of the charge incompressible quantum liquid state of an interacting charge  $2e$  bose fluid in 2d, where the insulating phase can be thought of as a vortex pair condensed state. This could exhibit QNF with a fractional charge  $e^*(= \frac{2e}{2})$ . This is a non trivial quantum fluid, where the powerful charge-vortex duality can be invoked for further understanding. In view of the gap, the solitonic objects have induced charges that are 'localized' - in other words they have a finite size. In this situation the exchange statistics makes sense. And one can use relations such as  $\Phi_c \Phi_v = 1$  (analogue of Dirac's quantization condition derived using spatial exchanges) without any problem and this restricts allowed fractional charges to rational values (because the vorticity is always an integer).

Experiments in cuprates show a metallic spin gap state crossing over to a (spin)gapless metallic non fermi liquid state as we increase doping. Hence a physically meaningful **zero temperature reference state** is a metallic state rather than an insulating state. This reference non fermi liquid state undergoes a superconducting transition at a  $T_c$ . We have to discuss the quantum number fractionization issue in this reference metallic state rather than a charge insulator at finite doping.

The 'extended' character of holon with a power law form factor (discussed earlier) makes the exchange statistics lose its precise meaning and also the corresponding relation  $\Phi_c \Phi_v = 1$ . This was shown by Haldane and Wu [18] in the context of a suggestion of Chao et al. [19] for fractional exchange statistics for quantized vortices in superfluid  $^4\text{He}$  film. Even though exchange statistics loses its meaning, a new possibility arises, namely Haldane's exclusion statistics [13]. This notion is meaningful in the gapless system and is dimension independent. In some approximate studies [20] it has been proposed that the spinons and holons have non trivial exclusion statistics in the 2d t-J model.

In view of the above discussion we conclude that the QNF of  $e^* = \frac{2e}{2}$  is an artificial restriction in the spin gap phase of the cuprates. The  $Z_2$  character of the spin sector leads to the interesting possibility of  $\pi$  flux condensation (i.e., a cross over to the spin gap phase) from the uniform RVB state [21]. It is very likely that the charge of the holon (and its exclusion statistics) will continuously vary from  $e^* = e$  as a function of doping. This is also a fundamental difference of cuprates from the well gapped quantum Hall system or other charge insulators.

In conclusion, if quantum number fractionization oc-

curs in cuprates leading to the experimentally seen non-fermi liquid phase and the d-wave superconducting state, the only viable mechanism seems to be through the RVB mechanism of spin charge separation, which comes out in a natural fashion in the RVB  $U(1)$  gauge theory. The variational/RVB mean field theory method combined with Gutzwiller projection [1,21], or the tomographic Luttinger liquid approach [2] are different ways of solving the same complex problem.

I wish to thank R. Shankar (Madras) for discussions.

- 
- [1] P.W. Anderson, Science **235** 1196 (87)
  - [2] P.W. Anderson, The theory of high temperature superconductivity (Princeton University Press, NY, 1996)
  - [3] Z. Zou and P.W. Anderson, Phys. Rev. **B 37** 627 (88)
  - [4] G. Baskaran and P.W. Anderson, Phys. Rev. **B37** 580 (88)
  - [5] P.B. Wiegman, Physica Scripta **T27** 160 (89)
  - [6] G. Baskaran, Physica Scripta, **T27** 53 (89); G. Baskaran and R. Shankar, J.Mod. Phys. Lett.**B2**, 1211 (1988)
  - [7] S. Kivelson, J. Sethna and D. Rokhsar, Phys. Rev. **B 38** 8865 (87)
  - [8] G. Baskaran, Yu Lu and E. Tosatti, Int. J. Mod. Phys. **B 1** 555 (1988).
  - [9] J.B. Marston, Phys. Rev. Lett., **61** 1914 (1988)
  - [10] P.B. Wiegman, Phys. Rev. Lett. **60** 821 (88); Z. G. Wen and A. Zee, Phys. Rev. Lett. **62** 2873 (89); Z. G. Wen, F. Wilczek and A. Zee, Phys. Rev. **B39** 11413 (89); L. Ioffe and V. I. Larkin, Phys. Rev. **B 38** 8988 (89); P.A. Lee and N. Nagaosa, Phys. Rev. **45** 966 (92); X.-G. Wen and P.A. Lee, Phys. Rev. Lett. **80** 2193 (98); P.A. Lee et al. Phys. Rev. **57** 6003 (98); Dung-Hai Lee, Phys. Rev. Lett. **84** 2494(00)
  - [11] T. Senthil, M.P.A. Fisher, cond-mat/9910224; L. Balents, M.P.A. Fisher and C. Nayak, Phys. Rev. **B 60** 1654 (99)
  - [12] T. Senthil, M.P.A. Fisher, cond-mat/0006500
  - [13] F.D.M. Haldane, Phys. Rev. Lett. **67** 937 (91)
  - [14] I. Affleck and J. B. Marston, Phys. Rev. **B 37** 3774 (88)
  - [15] G. Baskaran, Phys. Rev. Lett., **63** 2524 (1989); N. Read and S. Sachdev, Phys. Rev. Lett., **66** 1773 (91); S. Sachdev and N. Read, Int. J. Mod. Phys. **B5** 219 (91); X.G. Wen, Phys. Rev., **B 44** 2664 (91)
  - [16] D. S. Rokhsar and S. A. Kivelson, Phys. Rev. Lett. **61** 2376 (88)
  - [17] B. Sutherland, Phys. Rev. **38** 7192 (88)
  - [18] F.D.M. Haldane and Yong-Shi Wu, Phys. Rev. Lett. **55** 2887 (85)
  - [19] R. Y. Chao et al., Phys. Rev. Lett., **54** 1339 (85)
  - [20] F. Basco et al. J. Phys. Soc. Jap., **65** 687 (1996)
  - [21] G. Baskaran, Z. Zou and P.W. Anderson, Sol. St. Commn, **63** 873 (87)